

# oscode: fast solutions of oscillatory ODEs in cosmology

Fruzsina J. Agocs \* 1,2

<sup>1</sup>Kavli Institute for Cosmology, Cambridge

<sup>2</sup>Astrophysics Group, Cavendish Laboratory, Cambridge

**oscode** is a Python/C++ package for the fast solution of oscillatory ordinary differential equations. It handles equations of the form  $\ddot{x}(t) + 2\gamma(t)\dot{x}(t) + \omega^2(t)x(t) = 0$ . The time-dependence of the frequency  $\omega$  and the damping term  $\gamma$  can be explicit or implicit; below are examples of both. The algorithm is significantly more efficient than conventional (Runge-Kutta-based) solvers found in numerical libraries, thanks to reduced number of steps needed to traverse highly oscillatory regions.

## Airy equation




Figure 1: Steps and accuracy of **oscode** solving  $\ddot{x} + tx = 0$ .

- **Left:** different coloured dots show the internal steps **oscode** takes in 'RK' and 'WKB' mode, the continuous line being the exact solution
- **Right:** relative accuracy of **oscode**'s approximate solution
- Compare to scenario where **oscode** only takes 'RK' steps

## Burst equation




Figure 2: Solving  $\ddot{x} + \frac{n^2-1}{(1+t^2)^2}x = 0$ , an equation exhibiting a burst of oscillations.

- Equation has a parameter  $n$ ,  $n \sim$  no. of oscillations in total
- **Left:** **oscode**'s internal steps for  $n = 40$
- **Middle:** Relative error in **oscode**'s solution, and error in 'pure RK' solution for comparison
- **Right:** number of oscillations crossed per step for  $n = 10^5$

## Schrödinger equation




Figure 3: Eigenstates of the Schrödinger equation ( $\Psi'' + 2m(E - V(x))\Psi = 0$ ) in a harmonic potential well.

- Above is an example where the eigenenergies are known, but can compute them numerically:
- Start integration from well outside potential well on either side and meet at  $x = 0.5$
- Compute  $\Psi'/\Psi$  at the meeting point for both solutions; the difference is minimal if the energy is an eigenvalue
- Calculated energy eigenvalues in a harmonic well with quartic anharmonicity up to the 10000<sup>th</sup> energy level to 1 in  $10^7$  accuracy

## Primordial power spectra





Figure 4: Scalar primordial power spectra in closed universes with varying start-of-inflation curvature  $\Omega_k^i$ .

- Solve the Mukhanov-Sasaki (MS) equation for each wavenumber  $k$ , frequency is roughly proportional to  $k$
- MS equation is an example of an oscillator where  $\omega$  and  $\gamma$  are not explicit functions of time, are only available numerically
- **oscode** deals with this by performing interpolation on a grid of  $\omega$  and  $\gamma$  values
- Can test models where the perturbations have to be computed while they are deep inside the Hubble horizon and go through many oscillations before freeze-out.

## Algorithm

- **RK (Runge-Kutta) step:** approximates the solution at a later time  $t + h$  as a Taylor expansion around  $t$ .
- **WKB (Wentzel-Kramers-Brillouin) step:** uses an analytic approximation which gets better the slower the frequency of oscillations change, allowing **oscode** to traverse highly oscillatory regions in few steps.
- **oscode** switches between RK/WKB mode, choosing whichever allows for the largest step within a given error tolerance.
- The stepsize is updated to ensure the local error stays within the given tolerance.
- Derivatives of the frequency and damping term are required for both the RK and WKB steps, these are computed numerically as finite differences.
- WKB steps also require integrals, which are computed using Gauss-Lobatto integration.
- The RK steps and finite difference formulae were designed to minimise function evaluations.

For full paper, code and an animated version of Fig. 4, take a photo of this QR code or visit



<https://github.com/fruzsinaagocs/oscode>